

 HÉMA-QUÉBEC	Produits sanguins Cellules souches Tissus humains	Cord blood collection	PFN-00028[6]
		Effective date : 01-05-2023	Page 1 de 6

Table of contents

1.	Purpose	2
2.	Scope	2
3.	Responsibilities.....	2
4.	Required material.....	2
5.	Identifiers	2
6.	Procedure	3
6.1.	General.....	3
6.2.	Evaluation of the mother	4
6.3.	Umbilical cord disinfection	4
6.4.	Collection.....	5
7.	Appendices	6
8.	References	6
9.	List of modifications	6

Pour copie papier seulement : Le document original est approuvé avec signature numérique sous la responsabilité de l'AQ.

Tous droits réservés. Aucune partie du présent document ne peut être reproduite, conservée en mémoire ou transmise, sous quelque forme que ce soit ou par quelque moyen que ce soit, photocopie, enregistrement, procédés électroniques et mécaniques ou autres, sans la permission écrite préalable d'Héma-Québec.



Produits sanguins
Cellules souches
Tissus humains

Cord blood collection

PFN-00028[6]

Page 2 de 6

1. PURPOSE

To describe the cord blood collection procedure

2. SCOPE

All cord blood collections

3. RESPONSIBILITIES

Maternity ward and operating room physicians

4. REQUIRED MATERIAL

- Collection bag

5. IDENTIFIERS

N/A




6. PROCEDURE

6.1. GENERAL

- Regarding any questions, Hema-Quebec Public Cord Blood Bank can be reached at 1-888-666-4362 p.5253.
- While recruiting a potential cord blood donor, use the folder provided by Hema-Quebec that contains the PUB-00058 *Information guide on cord blood donation* (PUB-00057).
- The collection techniques must not jeopardize the health and safety of the mother and baby.
 - For mothers delivering at less than 34 weeks of gestation, the cord blood collection may be performed if the maternity ward physician considers it safe.
- A minimum of two collections per trained physician per year must be performed to remain qualified. Hema-Quebec will perform a quarterly follow-up to this effect.
- All materials required for the cord blood collection must be on a sterile field prior to use.
- The collection bag must only be manipulated aseptically.
- Collection may take a few minutes, and is performed after the delivery of the infant, and before delivery of the placenta.
- When handling the collection bag, pay special attention to the needle protector to prevent it from sliding against the needle.

Attention: Once the red clamp has been closed, it cannot be re-opened.



6.2. EVALUATION OF THE MOTHER

- Evaluate if the mother presents:
 - signs of infection such as fever (over 39°C);
 - evidence of parenteral drug use.
- If so, do not perform collection and document the reason in the « Commentaires / Écarts aux procédures » section of *Collecte de sang de cordon* form.

6.3. UMBILICAL CORD DISINFECTION

- After the birth of the baby, place a sterile field on the mother's perineal area and place the umbilical cord on this field.
- Open the plastic bag containing the collection bag and remove only the tubing.
- Visually inspect the collection bag and the tubing to ensure it is in good condition.
 - Return all defective material to Hema-Quebec.
- Close the blue clamps.
- Disinfect the cord unidirectionally.

6.4. COLLECTION

- Insert needle into the umbilical vein.



- Open the blue clamps and lower the collection bag to allow the cord blood to flow by gravity.

Note: A blood cord unit that is more than 85 mL will have better odds to attain qualification. However, all units collected must be sent to Hema-Quebec, without exception.

- After all the cord blood has been drained, milk the cord from the placenta towards the clamp.
- Close the blue clamps, then withdraw the needle from the vein.
- Push the needle protector up to make sure it covers the needle and lock it into position.
- Break the cannula from the anticoagulant pouch.
- Open the proximal blue clamp.
- Roll the anticoagulant pouch up onto itself to empty the anticoagulant into the collection line.
- Close the proximal blue clamp and the permanent red clamp.
- Hand the blood bag over to the nurse.
- In section 2 of the *Collecte de sang de cordon* form (or the *Collecte de Sang de Cordon Dirigé*), sign with your name and medical license number.
- In the event that the collection must be interrupted or if any problems arise during collection, document details in the « Commentaires / Écarts aux procédures » section of *Collecte de sang de cordon* form.

 HÉMA-QUÉBEC	Produits sanguins Cellules souches Tissus humains	Cord blood collection	PFN-00028[6]
Page 6 de 6			

7. APPENDICES

N/A

8. REFERENCES

- Cord blood collection bag REF MSC1200PU MacoPharma NOTEURO121 (Rev03_20190611 and Rev04_20201021)

9. LIST OF MODIFICATIONS

SECTION	MODIFICATIONS	JUSTIFICATION	ORIGIN OF INFORMATION
3	Addition of « and operating room »	The procedure is applicable to physicians of the operating room that proceed to cord blood collection following a caesarean delivery	N/A
6.1 1 st arrow	Addition of contact information for Hema-Quebec Public Cord Blood Bank	NetCord-FACT standard	NetCord-FACT Standard B3.1.6
6.1 2 nd arrow	Addition of the documentation that is used in the recruitment process of a potential cord blood donor.	Audit Citation NetCord-FACT 2022	AUD-R00121 / AEX-2806-EXP
6.4 2 nd arrow	Addition of a note « A blood cord unit that is more than 85 mL will have better odds to attain qualification. However, all units collected must be sent to Hema-Quebec, without exception. »	Give information regarding the volume of cord blood to be collected.	N/A
References	Addition of insert revision Rev04_20201021	Update of current version	ENR-02840[5]

Reference: OC-08148